

Radiative corrections to deeply virtual Compton scattering^{*)}

A.V. BELITSKY^a, D. MÜLLER^b, L. NIEDERMEIER^b, A. SCHÄFER^b

^a*C.N. Yang Institute for Theoretical Physics,
State University of New York at Stony Brook
NY 11794-3800, Stony Brook, USA*

^b*Institut für Theoretische Physik, Universität Regensburg
D-93040 Regensburg, Germany*

Received XXX

We discuss possibilities of measurement of deeply virtual Compton scattering amplitudes via different asymmetries in order to access the underlying skewed parton distributions. Perturbative one-loop coefficient functions and two-loop evolution kernels, calculated recently by a tentative use of residual conformal symmetry of QCD, are used for a model dependent numerical estimation of scattering amplitudes.

1 Introduction.

The asymptotic freedom of QCD allows to calculate short distance quantities perturbatively from the first principles. Together with the use of factorization theorems, the cross sections for (semi-)inclusive reactions are predicted at a large scale as a convolution of perturbative parton cross sections and long distance parton distributions. In the case of exclusive reactions, such as two-photon reactions in the light-cone dominated region or diffractive processes, the amplitudes factorizes in hard-scattering amplitudes and hadron distribution amplitudes and/or skewed parton distributions (SPD). Such distributions were already proposed in the past [1] and they become more popular nowadays [2] since it has been realized their vital rôle for the proper description of the mentioned reactions. Moreover, as it was pointed out by J. Xi, their measurement can clarify the issue of how the hadron total angular momentum is distributed among its constituents. One can expect that the study of the SPDs will be one of the central issues of hadronic physics in nearest future.

A promising way to access the SPDs is to probe the nucleon with a highly virtual photon via the Compton scattering process, which is called deeply virtual Compton scattering (DVCS). Indeed, it is auspicious that a first evidence of experimental feasibility to detect such a signal has been revealed by the ZEUS collaboration [3]. On the theoretical side the factorization of the collinear singularities has been proved at leading twist-two level to all orders in coupling [4]. The local version of the OPE does not converge in the considered kinematical region. Fortunately, this problem can be overcome by a non-local light-cone expansion in terms of light-ray operators [5, 1]. Moreover, from the renormalization group equation of these operators one can derive in a straightforward manner the evolution equations for SPDs.

^{*)} Talk given by D. Müller at Inter. Workshop "PRAHA-Spin99", Prague, Sept. 6-11, 1999.

Here we shortly review our approach to complete next-to-leading (NLO) calculation, which is accomplished by a heavy use of the conformal symmetry to map the known results from forward deep inelastic scattering (DIS) to the non-forward kinematics of DVCS and simple one-loop addenda. The problems within our modus operandi have been resolved by employing conformal Ward identities and constraints arising from the algebra of the collinear conformal group [6].

2 The virtual Compton scattering process.

The virtual Compton scattering (VCS) process on a nucleon can be measured in the reaction

$$\ell^\pm N \rightarrow \ell^\pm N \gamma. \quad (1)$$

However, as shown in Fig. 1, there is contamination from the so-called Bethe-Heitler (BH) process in which the final real photon is radiated from the lepton beam. Thus,




Fig. 1. The virtual Compton scattering process (a) and the Bethe-Heitler process (b).

the scattering amplitude square $|T|^2$ in the cross section,

$$d\sigma = \frac{1}{4kP_1} |T|^2 (2\pi)^4 \delta^4(k + P_1 - k' - P_2 - q_2) \frac{d^3 \vec{k}'}{2\omega' (2\pi)^3} \frac{d^3 \vec{P}_2}{2E_2 (2\pi)^3} \frac{d^3 \vec{q}_2}{2\nu (2\pi)^3}, \quad (2)$$

contains beside the VCS and BH parts also the interference term:

$$|T|^2 = \sum_{\lambda', S_2, \Lambda} \{ |T_{VCS}|^2 + |T_{BH}|^2 + T_{VCS} T_{BH}^* + T_{VCS}^* T_{BH} \}. \quad (3)$$

The BH-amplitude is purely real and is given as a contraction of the leptonic tensor

$$L_{\mu\nu} = \bar{u}(k', \lambda') [\gamma_\mu (\not{k} - \not{\Delta})^{-1} \gamma_\nu + \gamma_\nu (\not{k}' + \not{\Delta})^{-1} \gamma_\mu] u(k, \lambda) \quad (4)$$

with the hadronic current $J_\nu = \bar{u}(P_2) \{ F_1(\Delta^2) \gamma_\nu + i F_2(\Delta^2) \sigma_{\nu\tau} \frac{\Delta^\tau}{2M} \} u(P_1)$:

$$T_{BH} = -\frac{e^3}{\Delta^2} \epsilon^*(\Lambda)_\mu L^{\mu\nu} J_\nu, \quad \text{where } \Delta = P_1 - P_2. \quad (5)$$

Since Dirac and Pauli form factors F_1 and F_2 are (partially) known from experimental measurements and can be parametrized to a certain accuracy by double dipole formulae, one can disentangle the content of the hadronic tensor T_{VCS} :

$$T_{VCS} = \mp \frac{e^3}{q_1^2} \epsilon^*(\Lambda)_\mu T^{\mu\nu} \bar{u}(k') \gamma_\nu u(k), \text{ where } \begin{cases} - & \text{for } e^- \\ + & \text{for } e^+ \end{cases}. \quad (6)$$

This tensor, defined by the time ordered product of two electromagnetic currents

$$T_{\mu\nu}(P, q_1, q_2) = i \int dx e^{ix(q_1+q_2)} \langle P_2, S_2 | T \{ j_\mu(x/2) j_\nu(-x/2) \} | P_1, S_1 \rangle, \quad (7)$$

contains for a spin-1/2 target twelve¹⁾ independent kinematical structures [7].

Measuring the cross section (2) in quite different experiments: with negatively or positively charged lepton beam, with (un)polarized beam or (un)polarized target, the rich structure of the hadronic tensor can be explored in different kinematical regions. In the following we are interested in DVCS, where $-q_1^2$ is large and Δ^2 is small. Unfortunately, since the DVCS amplitude (6) decreases with increasing q_1^2 while the BH amplitude (5) grows up with decreasing Δ^2 , in this kinematics it will be a delicate task to extract the VCS amplitude for large virtualities $-q_1^2$ and especially low Δ^2 from experimental data. On the other hand it is promising to look for asymmetries which allow one to measure the interference term directly and get in this way access to the real and imaginary part of the DVCS amplitude [8]:

1. charge asymmetry

$$d\sigma(e^- \text{-beam}) - d\sigma(e^+ \text{-beam}), \propto 4\text{Re}T_{DVCS}T_{BH} \quad (8)$$

2. single spin asymmetries (either the lepton-beam or the nucleon is polarized)

$$d\sigma(\uparrow) - d\sigma(\downarrow) \propto 4\text{Im}T_{DVCS}T_{BH}, \quad (9)$$

3. azimuthal angle asymmetry

$$A = \frac{\int_{-\pi/2}^{\pi/2} d\phi_r d\sigma - \int_{\pi/2}^{3\pi/2} d\phi_r d\sigma}{\int_0^{2\pi} d\phi_r d\sigma}, \quad (10)$$

where $\phi_r = \phi_N - \phi_e$ with $\phi_N(\phi_e)$ is the azimuthal angle of the final nucleon (lepton).

¹⁾ 12 = $\frac{1}{2} \times 3$ (virtual photon) $\times 2$ (final photon) $\times 2$ (initial nucleon) $\times 2$ (final nucleon). The reduction factor 1/2 is a result of parity invariance.

3 Practical implication of conformal symmetry.

To describe different “two-photon” processes in the light-cone dominated region in a unique way, we generalize the Bjorken limit and introduce the scaling variables:

$$\xi \equiv \frac{Q^2}{Pq}, \quad \eta \equiv \frac{\Delta q}{Pq}, \quad Q^2 = -q^2 = -\frac{1}{4}(q_1 + q_2)^2, \quad P = P_1 + P_2, \quad (11)$$

where $\xi \approx x_{\text{Bj}}/(2 - x_{\text{Bj}})$ is related to the Bjorken variable $x_{\text{Bj}} = -q_1^2/2(P_1 q_1)$ and η is the so-called skewedness parameter. Formally, this parameter interpolates between different processes as given in table 1.

DIS	$\Delta = 0$	$\eta = 0$
DVCS	$q_2^2 = 0$	$\eta = 1/\omega$
$\gamma^* N \rightarrow N l^+ l^-$	$q_2^2 > 0$	$\eta \approx \cos \phi_{\text{Br}} = \mathbf{p}_1 \mathbf{Q} / \mathbf{p}_1 \mathbf{Q} $, Breit-frame
$\gamma^* \gamma^* \rightarrow H H$		$\eta \approx \cos \phi_{\text{cm}} = \mathbf{p}_1 \mathbf{Q} / \mathbf{p}_1 \mathbf{Q} $, center-of-mass frame
$\gamma^* \gamma^* \rightarrow M$	$p_1 = 0$	$\eta = 1$

Table 1. The value of the scaling variable η for different two-photon processes.

Since in the case of DVCS the longitudinal part of the hadronic tensor vanishes, there are only two kinematical structures left to leading twist-two accuracy:

$$T_{\mu\nu} = -\tilde{g}_{\mu\nu}^T \mathcal{F}_1(\xi, \eta = \xi, Q^2, \Delta^2) + i\tilde{\epsilon}_{\mu\nu q P} \frac{1}{Pq} \mathcal{G}_1(\xi, \eta = \xi, Q^2, \Delta^2) + \dots \quad (12)$$

Here the transverse part of the metric tensor, denoted by $g_{\mu\nu}^T$, and the ϵ -tensor are contracted with the projection operators $\mathcal{P}_{\alpha\beta} = g_{\alpha\beta} - q_{1\alpha} q_{2\beta} / (q_1 q_2)$: $\tilde{g}_{\mu\nu}^T = \mathcal{P}_\mu^\alpha g_{\alpha\beta}^T \mathcal{P}_\nu^\beta$, $\tilde{\epsilon}_{\mu\nu q P} = \mathcal{P}_\mu^\alpha \epsilon_{\alpha\beta q P} \mathcal{P}_\nu^\beta$. The appearing amplitudes can be calculated by means of the OPE. In contrast to the case of DIS, a larger number of (local) operators and their Wilson coefficients contribute in the non-forward case. Fortunately, we can appreciate the conformal OPE (COPE) [9], which tells us that all Wilson coefficients are fixed up to an overall normalization, which can be taken from known results in DIS. The conformal operators appearing in the COPE are:

$$\begin{aligned} \left\{ \begin{matrix} Q\mathcal{O}^V \\ Q\mathcal{O}^A \end{matrix} \right\}_{jl} &= \bar{\psi}(i\partial_+)^l \left\{ \begin{matrix} \gamma_+ \\ \gamma_+ \gamma_5 \end{matrix} \right\} C_j^{\frac{3}{2}} \left(\frac{\overleftrightarrow{D}_+}{\partial_+} \right) \psi, \\ \left\{ \begin{matrix} G\mathcal{O}^V \\ G\mathcal{O}^A \end{matrix} \right\}_{jl} &= G_{+\mu} (i\partial_+)^{l-1} \left\{ \begin{matrix} g_{\mu\nu} \\ i\epsilon_{\mu\nu-+} \end{matrix} \right\} C_{j-1}^{\frac{5}{2}} \left(\frac{\overleftrightarrow{D}_+}{\partial_+} \right) G_{\nu+}, \end{aligned} \quad (13)$$

where $\partial = \overrightarrow{\partial} + \overleftarrow{\partial}$ and $\overleftrightarrow{D} = \overrightarrow{D} - \overleftarrow{D}$. The $+$ and $-$ components are obtained by contraction with the two light-like vectors n and n^* , such that $n^2 = n^{*2} = 0$ and $nn^* = 1$. They carry the spin quantum number $l+1$ and the conformal spin $j+1$.

In the conformal subtraction (CS) scheme their mixing is only induced by the trace anomaly and proportional to β [10]:

$$\mu \frac{d}{d\mu} \mathcal{O}_{jl} = \gamma_{jj} \mathcal{O}_{jl} + \frac{\beta}{g} \sum_{k=0}^{j-2} \Delta_{jk} \mathcal{O}_{kl} \quad (14)$$

The anomalous dimension matrix γ_{jj} is, up to a normalization factor in the mixed channels, the same as in DIS.

Unfortunately, for the case of DVCS kinematics the COPE in its local version does not converge. Thus, it is necessary to resum the local operators into so-called light-ray ones. This provide us a definition for the SPDs:

$$\begin{aligned} \left\{ \begin{matrix} Q_q^V \\ Q_q^A \end{matrix} \right\} (t, \eta) &= \int \frac{d\kappa}{2\pi} e^{i\kappa t P_+} \langle P_2 S_2 | \bar{\psi}(-\kappa n) \left\{ \begin{matrix} \gamma_+ \\ \gamma_+ \gamma_5 \end{matrix} \right\} \psi(\kappa n) | P_1 S_1 \rangle, \\ \left\{ \begin{matrix} G_q^V \\ G_q^A \end{matrix} \right\} (t, \eta) &= \frac{4}{P_+} \int \frac{d\kappa}{2\pi} e^{i\kappa t P_+} \langle P_2 S_2 | G_{+\mu}^a(-\kappa n) \left\{ \begin{matrix} g_{\mu\nu} \\ i\epsilon_{\mu\nu-+} \end{matrix} \right\} G_{\nu+}^a(\kappa n) | P_1 S_1 \rangle. \end{aligned} \quad (15)$$

The momentum fractions of the incoming [outgoing] parton is $\frac{t+\eta}{1+\eta} P_+$ [$\frac{t-\eta}{1+\eta} P_+$]. A form factor decomposition would give us the functions H , E and $\tilde{H}$, $\tilde{E}$ for the parity even and odd sectors, respectively, as introduced by Ji [2].

Next we have to reconstruct the coefficient functions appearing in

$$\mathcal{T}^a = \sum_Q e_Q^2 \left[Q T^a \otimes Q q^a + \frac{1}{N_f} \frac{1}{\eta} G T^a \otimes G q^a \right], \quad (16)$$

where $\mathcal{T}^V = \mathcal{F}_1$, $\mathcal{T}^A = \mathcal{G}_1$ and the sum runs over quark species $Q = u, d, s$ with electrical charge e_Q . The convolution is defined by $\tau_1 \otimes \tau_2 \equiv \tau_1(x, y) \tau_2(y, z)$. This task can be achieved by some manipulation of the COPE and, e.g. for the flavour non-singlet case, can be cast into the form

$$T(\xi, \eta, t, Q^2, \mu^2) = \frac{1}{\xi} F \left(\frac{\xi}{\eta}, \frac{r}{\eta} \right) \otimes \left(\frac{Q^2}{\mu^2} \right)^{V(\frac{r}{\eta}, \frac{s}{\eta})} \otimes C \left(\frac{s}{\eta}, \frac{t}{\eta}; \alpha_s \right). \quad (17)$$

The evolution kernel $V \left(\frac{r}{\eta}, \frac{s}{\eta} \right)$ and the coefficient function $C \left(\frac{s}{\eta}, \frac{t}{\eta}; \alpha_s \right)$ are diagonal w.r.t. the conformal waves — the Gegenbauer polynomials:

$$\eta^j C_j^{3/2} \left(\frac{t}{\eta} \right) \otimes \left\{ \begin{matrix} V \\ C \end{matrix} \right\} \left(\frac{t}{\eta}, \frac{t'}{\eta}; \alpha_s \right) = \left\{ \begin{matrix} -\frac{1}{2} \gamma_j(\alpha_s) \\ c_j(\alpha_s) \end{matrix} \right\} \eta^j C_j^{3/2} \left(\frac{t'}{\eta} \right), \quad (18)$$

and their eigenvalues $\gamma_j(\alpha_s)$ and $c_j(\alpha_s)$ coincide with anomalous dimensions and Wilson coefficients, respectively, which appear in DIS. A closed form of the function

F can be deduced order by order in α_s from the sum

$$F(x, y) = \sum_{j=0}^{\infty} \left(\frac{2x}{1+x} \right)^{j+1} \frac{B(j+1, j+2)}{(1+x)^{\gamma_j/2}} \times {}_2F_1 \left(\begin{matrix} 1+j+\gamma_j/2, 2+j+\gamma_j/2 \\ 4+2j+\gamma_j \end{matrix} \middle| \frac{2x}{1+x} \right) C_j^{3/2}(y). \quad (19)$$

Conformal covariance does not hold true in the minimal subtraction (MS) scheme. The transformation, which rotates the CS result (17-19) to the minimal one, is induced by the conformal anomalies appearing in the MS scheme. The rotation matrix can be found by study of the conformal Ward identities for Green functions with operator insertions and calculating the anomalous contributions via (modified) Feynman rules. Indeed, the off-diagonal piece of the anomalous dimension matrix to NLO is induced by the special conformal anomaly matrix $\gamma^{(0)c}$ and the first coefficient of the β -function [6]:

$$\gamma_{jk}^{(1)} = \gamma_{jj}^{(1)} - \frac{[\gamma^{(0)c} + \beta_0 \mathbf{b}, \gamma^{(0)}]_{jk}}{2(j-k)(j+k+3)}, \quad \mathbf{b}_{jk} = -[1 + (-1)^{j-k}] \theta_{jk}(2k+3) \mathbf{1}. \quad (20)$$

The remaining task to transform the coefficient functions to the MS scheme have been solved already in [10]. In the case of DVCS, they are nothing else as the analytical continuation of the hard-scattering amplitudes for the meson production by two photon-collision. In the case of the axial-vector channel they read:

$$\begin{aligned} Q_T^{(0)} &= \frac{1}{1-t}, \quad G_T^{(0)} = 0, \\ Q_T^{A(1)} &= \frac{C_F}{2(1-t)} \left[\left(2 \ln \frac{1-t}{2} + 3 \right) \left(\frac{1}{2} \ln \frac{1-t}{2} - \frac{3}{4} \right) - \frac{27}{4} - \frac{1-t}{1+t} \ln \frac{1-t}{2} \right], \\ G_T^{A(1)} &= \frac{N_f}{2} \left[\left(\frac{1}{1-t^2} + \frac{\ln \frac{1-t}{2}}{(1+t)^2} \right) \left(\ln \frac{1-t}{2} - 2 \right) - \frac{\ln^2 \frac{1-t}{2}}{2(1+t)^2} \right]. \end{aligned} \quad (21)$$

4 Numerical estimations.

There are different proposals for modeling the SPD. Here we will deal only with two of them, equating the SPD equal to the forward parton distributions (FPD-model) and using the proposal of Radyushkin for the ansatz of the double distribution (DD) functions [11] $f(z_-, z_+) = \pi(z_-, z_+) f(z_+)$, where the profile function π for quarks and gluons are given by

$$Q\pi(z_-, z_+) = \frac{3}{4} \frac{[1 - |z_+|^2] - z_-^2}{[1 - |z_+|^2]^3}, \quad G\pi(z_-, z_+) = \frac{15}{16} \frac{\{[1 - |z_+|^2] - z_-^2\}^2}{[1 - |z_+|^2]^5}, \quad (22)$$

The SPD is obtained by an integration [1]:

$$q(t, \eta, Q^2) = \int_{-1}^1 dz_+ \int_{-1+|z_+|}^{1-|z_+|} dz_- \delta(z_+ + \eta z_- - t) f(z_-, z_+, Q^2). \quad (23)$$

We apply the DD model at different scales. To evolve the SPD, we expand them:

$$q(t, \eta, Q^2) = \sum_{j=0}^{\infty} c_j(\eta, Q^2) P_j(t) \quad (24)$$

in terms of Legendre polynomials and express the expansion coefficients $c_j(\eta, Q^2)$ in terms of conformal moments which evolution is governed by the anomalous dimension matrix (20). This method is suited for not too small η .




Fig. 2. In (a) and (b) we show the SPD for the sum of the non-polarized u and $\bar{u}$ quarks and the gluon densities, respectively. The FPD-model (dotted line) and the DD-model with MRS [13] parametrization (dashed line) are modeled at a scale of 4 GeV^2 , while the DD-Model with GRV [14] parametrization (solid line) has been taken at the momentum scale 0.4 GeV^2 and has been evolved afterwards with NLO formulae to $Q_0^2 = 4 \text{ GeV}^2$. The skewedness parameter is $\eta = 1/3$ which corresponds to $x_{\text{Bj}} = 1/2$.

In Fig. 2 we show our input distributions at a scale of $Q^2 = -q_1^2 = 4 \text{ GeV}^2$. Let us also mention, that the relative NLO corrections due to the evolution are as expected not very large. Starting at an input scale of 0.7 GeV , we found that the NLO corrections do not exceed 10% to 30% [12].

For the numerical estimation of the hard-scattering amplitudes (16) we perform also an expansion with respect to Legendre polynomials which allows us to include easily evolution effects. However, this method is not appropriate to study the low x_{Bj} region. In general we found that the NLO corrections for $x_{\text{Bj}} > 0.1$ are typically of the order of 20-50% (see Fig. 3a). In special cases, e.g. for the DD model with MRS parameterization, it can even be much larger. Fortunately, there is partial




Fig. 3. The real part (a) of $\mathcal{F}_1$ for DD-model with MRS and GRV parametrizations for the scale $Q_0^2 = 4 \text{ GeV}^2$ and for $x_{Bj} \geq 0.1$. Here MRS (GRV) amplitude is plotted at LO as dashed (dotted) and at NLO as solid (dash-dotted) lines, respectively. The ratio of the real to imaginary part (b) of $\mathcal{F}_1$, respectively, for the DD- and FPD-model with the MRS parametrization. Here the DD(FPD)-model at LO and NLO is shown as dashed (dotted) and solid (dash-dotted) lines, respectively. For the SPD-models considered the result is independent of Δ^2 .

cancellation of perturbative corrections in the ratio of real to imaginary part. As shown in Fig. 3b, this ratio is model dependent and may its measurement can be used to get a deeper insight in the SPD.

A.B. was supported by the Alexander von Humboldt foundation.

References

- [1] D. Müller et al., Fortschr. Phys. 42 (1994) 101.
- [2] X. Ji, Phys. Rev. D 55 (1997) 7114; A.V. Radyushkin, Phys. Lett. B 380 (1997) 417; J.C. Collins, L. Frankfurt, M. Strikman, Phys. Rev. D 56 (1997) 2982.
- [3] P.R.B. Saul (for ZEUS Collaboration), Prompt photon production and deeply virtual Compton scattering, talk at EPS99, Tampere, Finland (July, 1999).
- [4] A.V. Radyushkin, Phys. Rev. D 56 (1997) 5524; J.C. Collins, A. Freund, Phys. Rev. D 59 (1999) 074009; X. Ji, J. Osborne, Phys. Rev. D 58 (1998) 094018.
- [5] S.A. Anikin, O.I. Zavialov, Ann. Phys. 116 (1978) 135; B. Geyer et al., Z. Phys. C 26 (1985) 591; I.I. Balitsky, V.M. Braun, Nucl. Phys. B 311 (1989) 541.
- [6] D. Müller, Phys. Rev. D 49 (1994) 2525; A.V. Belitsky, D. Müller, Nucl. Phys. B 527 (1998) 207, Nucl. Phys. B 537 (1999) 397.
- [7] P. Kroll, M. Schürmann, P.A.M. Guichon, Nucl. Phys. A 598 (1996) 435.
- [8] L.L. Frankfurt, A. Freund, M. Strikman, Phys. Rev. D 58 (1998) 114001; P.M. Guichon, M. Vanderhaeghen, Prog. Part. Nucl. Phys. 41 (1998) 125; A. Freund, M. Strikman, Single Spin Asymmetry in DVCS, hep-ph/9906205.

- [9] S. Ferrara, R. Gatto, A.F. Grillo, Phys. Lett. B 36 (1971) 124, Phys. Rev. D 5 (1972) 3102.
- [10] D. Müller, Phys. Rev. D 58 (1998) 054005; A.V. Belitsky, D. Müller, Phys. Lett. B 417 (1997) 129.
- [11] I.V. Musatov and A.V. Radyushkin, Evolution and Models for Skewed Parton Distributions, hep-ph/9905376.
- [12] A.V. Belitsky, et al., Phys. Lett. B 437 (1998) 160, Nucl. Phys. B 546 (1999) 279.
- [13] A.D. Martin, R.G. Roberts, W.J. Stirling, Phys. Rev. D 47 (1993) 867.
- [14] M. Glück, E. Reya, A. Vogt, Eur. Phys. J. C 5 (1998) 461.